

# Global, regional, and national burden of household air pollution, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021



GBD 2021 HAP Collaborators\*

## Summary

**Background** Despite a substantial reduction in the use of solid fuels for cooking worldwide, exposure to household air pollution (HAP) remains a leading global risk factor, contributing considerably to the burden of disease. We present a comprehensive analysis of spatial patterns and temporal trends in exposure and attributable disease from 1990 to 2021, featuring substantial methodological updates compared with previous iterations of the Global Burden of Diseases, Injuries, and Risk Factors Study, including improved exposure estimations accounting for specific fuel types.

**Methods** We estimated HAP exposure and trends and attributable burden for cataract, chronic obstructive pulmonary disease, ischaemic heart disease, lower respiratory infections, tracheal cancer, bronchus cancer, lung cancer, stroke, type 2 diabetes, and causes mediated via adverse reproductive outcomes for 204 countries and territories from 1990 to 2021. We first estimated the mean fuel type-specific concentrations (in  $\mu\text{g}/\text{m}^3$ ) of fine particulate matter ( $\text{PM}_{2.5}$ ) pollution to which individuals using solid fuels for cooking were exposed, categorised by fuel type, location, year, age, and sex. Using a systematic review of the epidemiological literature and a newly developed meta-regression tool (meta-regression: Bayesian, regularised, trimmed), we derived disease-specific, non-parametric exposure–response curves to estimate relative risk as a function of  $\text{PM}_{2.5}$  concentration. We combined our exposure estimates and relative risks to estimate population attributable fractions and attributable burden for each cause by sex, age, location, and year.

**Findings** In 2021, 2.67 billion (95% uncertainty interval [UI] 2.63–2.71) people, 33.8% (95% UI 33.2–34.3) of the global population, were exposed to HAP from all sources at a mean concentration of  $84.2 \mu\text{g}/\text{m}^3$ . Although these figures show a notable reduction in the percentage of the global population exposed in 1990 (56.7%, 56.4–57.1), in absolute terms, there has been only a decline of 0.35 billion (10%) from the 3.02 billion people exposed to HAP in 1990. In 2021, 111 million (95% UI 75.1–164) global disability-adjusted life-years (DALYs) were attributable to HAP, accounting for 3.9% (95% UI 2.6–5.7) of all DALYs. The rate of global, HAP-attributable DALYs in 2021 was 1500.3 (95% UI 1028.4–2195.6) age-standardised DALYs per 100000 population, a decline of 63.8% since 1990, when HAP-attributable DALYs comprised 4147.7 (3101.4–5104.6) age-standardised DALYs per 100000 population. HAP-attributable burden remained highest in sub-Saharan Africa and south Asia, with 4044.1 (3103.4–5219.7) and 3213.5 (2165.4–4409.4) age-standardised DALYs per 100000 population, respectively. The rate of HAP-attributable DALYs was higher for males (1530.5, 1023.4–2263.6) than for females (1318.5, 866.1–1977.2). Approximately one-third of the HAP-attributable burden (518.1, 410.1–641.7) was mediated via short gestation and low birthweight. Decomposition of trends and drivers behind changes in the HAP-attributable burden highlighted that declines in exposures were counteracted by population growth in most regions of the world, especially sub-Saharan Africa.

**Interpretation** Although the burden attributable to HAP has decreased considerably, HAP remains a substantial risk factor, especially in sub-Saharan Africa and south Asia. Our comprehensive estimates of HAP exposure and attributable burden offer a robust and reliable resource for health policy makers and practitioners to precisely target and tailor health interventions. Given the persistent and substantial impact of HAP in many regions and countries, it is imperative to accelerate efforts to transition under-resourced communities to cleaner household energy sources. Such initiatives are crucial for mitigating health risks and promoting sustainable development, ultimately improving the quality of life and health outcomes for millions of people.

**Funding** Bill & Melinda Gates Foundation.

**Copyright** © 2025 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY license.

## Introduction

Household air pollution (HAP) from solid cooking fuels is a known source of health-relevant exposure for about

3 billion people worldwide.<sup>1–3</sup> People living in households using primarily solid fuels (coal or charcoal, wood, crop residues, and dung) for cooking are exposed to high

Published Online  
March 18, 2025  
[https://doi.org/10.1016/S0140-6736\(24\)02840-X](https://doi.org/10.1016/S0140-6736(24)02840-X)  
See Online/Comment  
[https://doi.org/10.1016/S0140-6736\(25\)00467-2](https://doi.org/10.1016/S0140-6736(25)00467-2)

\*Collaborators listed at the end of the Article

Correspondence to:  
Dr Katrin G Burkart, Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA 98105, USA  
[katburk@uw.edu](mailto:katburk@uw.edu)

**Research in context****Evidence before this study**

Previous research on the global burden and mortality from household air pollution (HAP) used integrated exposure–response curves that included few HAP studies and relied on passive and active smoking data or included pooled relative risks that assumed individuals to be either exposed or unexposed to HAP. Such estimations also assumed exposure to equal concentrations of particulate matter with a diameter of less than 2.5 µm (PM<sub>2.5</sub>), regardless of location and fuel type used.

**Added value of this study**

This study extends previous efforts to estimate the burden attributable to HAP, with year-specific and location-specific exposures modelled by fuel type, focusing on coal, crop residues, dung, and wood. This innovative approach enabled a more precise estimation of exposure to PM<sub>2.5</sub> and the associated burden of disease than previous studies, allowing us to provide comprehensive and timely estimates for 204 locations from 1990 to 2021. Increased availability of epidemiological studies and a newly developed meta-regression tool (meta-regression: Bayesian, regularised, trimmed) allowed us to expand the scope of health outcomes and discontinue the use of active and second-hand smoking data in our exposure–response estimations. Notably, we present the first global analysis to incorporate causes mediated

via adverse reproductive outcomes—ie, short gestation and low birthweight. The indirect effect of HAP on reproductive health extends across the lifespan due to negative impacts on outcomes including diarrhoeal disease, respiratory disease, and other infectious and non-infectious diseases, substantially adding to the burden of disease. We also included cataract (estimated as a HAP-attributable cause for females since the Global Burden of Diseases, Injuries, and Risk Factors Study [GBD] 2010 and for males since GBD 2019) and diabetes (first included as a HAP-attributable cause in GBD 2017) in our estimations.

**Implications of all the available evidence**

Despite substantial efforts to reduce exposure and decreases in the proportion of individuals cooking with solid fuels, exposure to and burden from HAP remain high. Our findings highlight higher exposure, morbidity, and mortality than previously estimated and show that although reductions in exposure have led to a net decrease in HAP burden over time, population growth has counteracted much of the effect of reduced exposure. We found that about a third of HAP-attributable burden is mediated through adverse reproductive outcomes, leading to tens of millions of years of life lost annually. Renewed efforts and international investments are required to transition exposed communities to cleaner fuels and reduce the consequent lifelong burden of household air pollution.

levels of HAP from particulate matter with a diameter of less than 2.5 µm (PM<sub>2.5</sub>), a well-established hazard to human health.<sup>4</sup> Burns are common in households using solid fuels, and fuel collection typically falls to women and girls, consuming tens of hours per week.<sup>5</sup> UN Sustainable Development Goal (SDG) 3.9 aims to reduce morbidity and mortality from environmental pollution, and SDG 7.1 calls for universal access to cleaner fuels by 2030, but many countries still do not have the resources to achieve these goals.<sup>6</sup>

Previous studies, including the Global Burden of Diseases, Injuries, and Risk Factors Study, have estimated the HAP-attributable burden of disease, but limitations relating to methods and data remain. Past exposure estimates either did not account for spatiotemporal variability<sup>2,3</sup> or did not adjust household exposure to the individual level.<sup>7</sup> Previous estimates of burden relied on a binary exposure indicator<sup>2,4,8</sup> and did not include type 2 diabetes or causes mediated via adverse reproductive outcomes.<sup>8</sup> Here, we provide an improved methodological framework developed over the past decade that more accurately characterises exposure and risk estimates. Although previous studies assumed that different solid fuel types produce the same amount of PM<sub>2.5</sub> pollution,<sup>2,8</sup> a known simplification, in this study we modelled fuel type-specific exposure for four distinct categories—ie, crop residues, dung, wood, and coal and

charcoal—resulting in more precise PM<sub>2.5</sub> exposure estimates.<sup>9</sup> Moreover, we improved our exposure–response estimates by incorporating newly published studies. Thus, the meta-regression in this study does not include active and passive smoking studies, which previously served as substitutes for high exposures.<sup>10</sup> We also added type 2 diabetes and causes mediated via low birthweight and short gestation to our global analysis. This study is the first to detail the GBD methodology for estimating HAP-attributable burden since the assessment by Smith and colleagues<sup>3</sup> of the approach used for GBD 2010, and we present numerous methodological and input data updates.

On the basis of these methodological improvements and updated data regarding exposure and relative risk, we estimated the burden of disease attributable to HAP for 204 countries and territories from 1990 to 2021. In addition to analysing spatial patterns, we conducted decomposition analysis to better understand the drivers behind changes in the attributable burden and to evaluate the role of demographic changes versus changes in exposure to HAP over this 31-year period.<sup>11,12</sup> Our study adds to a growing body of work that shows the need for renewed action to reduce sustained burden from HAP. This manuscript was produced with input from the GBD Collaborator Network and in accordance with the GBD Protocol.

## Methods

### Overview

GBD 2021 complies with the GATHER statement (appendix 1 p 12).<sup>11</sup> Estimating the disease burden attributable to HAP required the following steps: global estimation of the number and proportion of people exposed to HAP by year, location, and fuel type; conversion of exposure proportions to fuel-type-specific  $PM_{2.5}$  concentrations by year, location, age, and sex; generation of exposure–response curves describing the relationships between  $PM_{2.5}$  concentration and relative risks (RRs) of incidence for specific diseases; and application of a theoretical minimum risk exposure level (TMREL)—ie, the level of risk exposure that minimises disease risk at the population level. These steps allowed for quantification of HAP-specific population attributable fractions that were then used to estimate HAP-attributable burden using GBD estimates for each year, age, sex, and location.

### Proportion of population exposed to solid fuel types

In the first step, we used survey data to estimate the proportion of the population primarily using solid fuels in each location and year. For the purposes of this study, solid fuel was specified as coal and charcoal, wood, crop residues, and dung (appendix 1 pp 4–10). Estimations were based on data from household surveys (eg, Demographic and Health Surveys and Multiple Indicator Cluster Surveys), population censuses, WHO's Household Energy Database,<sup>13</sup> and other sources, such as country-specific surveys. For the code used in this analysis, see appendix 1 (p 4); for sources and estimates, see appendix 2 (p 6). Count data are presented to three significant figures and rates (percentages) to one decimal place.

We used the GBD 2021 spatiotemporal Gaussian process regression tool, a three-stage non-linear model that allowed us to produce a full set of estimates of the proportion of the population using each solid fuel type for all years and locations (appendix 1 pp 4–9). The first stage of the modelling process is a linear regression informed by both exposure data and covariates, the second stage smooths data variation across locations and time by analysing the residuals, and the third stage uses Gaussian process regression to produce the final model fit. After modelling each fuel type and the overall solid fuel exposure categories, we adjusted the sum of the fuel types to fit the solid fuel category for each location and year.

### Mapping of exposure to $PM_{2.5}$

In the second step, we estimated the mean fuel type-specific concentration (in  $\mu\text{g}/\text{m}^3$ ) of  $PM_{2.5}$  pollution to which individuals using solid fuels for cooking were exposed, categorised by location, year, age, and sex (appendix 1 pp 10–14; for sources and estimates, see appendix 2 pp 4–5). This model enabled us to convert the

proportion of the population using a given solid fuel type to the excess concentration of  $PM_{2.5}$  to which people were exposed (above the ambient concentration) by age and sex using data from real-world observations of HAP concentrations.<sup>14–16</sup> Converting exposure proportions to fuel-type specific  $PM_{2.5}$  concentrations allowed us to use these data as direct input for  $PM_{2.5}$  risk curves.

See Online for appendix 1

### Modelling of RR

In the third step, we modelled HAP-attributable RR for the following outcomes: lower respiratory infection; stroke; ischaemic heart disease; chronic obstructive pulmonary disease (COPD); cancers of the trachea, bronchus, and lung; type 2 diabetes; cataract; low birthweight; and short gestation (appendix 1 pp 14–44; for sources, see appendix 2 p 6). Type 2 diabetes, low birthweight, and short gestation have been added as HAP-attributable causes in GBD analyses since GBD 2010.<sup>17,18</sup> As the adverse reproductive outcomes low birthweight and short gestation are already risk factors, we used a mediation analysis to reattribute the portion of the burden (ie, otitis, meningitis, encephalitis, sudden infant death syndrome, upper respiratory infections, diarrhoeal diseases, neonatal disorders, and lower respiratory infections) attributable to low birthweight and short gestation to  $PM_{2.5}$  pollution instead (appendix 1 pp 40–44).<sup>18</sup>

For each outcome except for cataract, we calculated a risk curve on the basis of epidemiological data characterising exposure to ambient  $PM_{2.5}$  air pollution and HAP. These curves enabled us to use the output of our mapping model to calculate the location-specific relative risk for a given cause. We used the recently developed burden of proof risk function framework incorporating a meta-regression tool—meta-regression: Bayesian, regularised, trimmed (MR-BRT)<sup>19,20</sup>—to generate flexible, non-linear RR splines (exposure–response curves), combine available RR studies while minimising the effects of outliers, characterise and correct systematic biases, account for between-study heterogeneity, adjust for confounders, and extrapolate beyond the exposure levels observed in most of the literature to the very high levels of pollution found in particular locations (appendix 1 pp 14–44). For cataract, we used the burden of proof risk function approach to calculate RR for exposed versus unexposed individuals (appendix 1 p 39).

See Online for appendix 2

The highest  $PM_{2.5}$  concentration attributed to HAP reported in the epidemiological literature to develop risk curves is approximately  $550 \mu\text{g}/\text{m}^3$ . However, because much higher  $PM_{2.5}$  concentrations have been documented in observational studies,<sup>14–16</sup> we extrapolated our risk curves to a concentration of  $1000 \mu\text{g}/\text{m}^3$ .

### Proportional population attributable fraction calculation and burden estimation


Population attributable fractions (PAFs) were estimated together for outcomes related to both ambient pollution

and HAP to account for combined exposure to both types of air pollution in a single location. The TMREL for ambient  $PM_{2.5}$  was defined as uniform distribution between  $2.4 \mu\text{g}/\text{m}^3$  and  $5.9 \mu\text{g}/\text{m}^3$   $PM_{2.5}$  (appendix 1 p 10) and, therefore, assumed no incremental HAP exposure.<sup>21</sup> Because cataract is a consequence of HAP only, not ambient particulate matter, unlike other outcomes, the TMREL for cataract was defined as individuals not using solid cooking fuel. The PAF quantifies the fraction of burden that could be attributed to a risk factor if the risk factor was reduced to the TMREL. To calculate PAFs, we first calculated the exposure to particulate matter as the sum of the ambient concentration of  $PM_{2.5}$  and household or indoor concentration of  $PM_{2.5}$ . We used the population-weighted mean ambient  $PM_{2.5}$  concentration for a specific location and year and calculated indoor  $PM_{2.5}$  concentrations as a function of the percentage of population exposed and the concentrations of  $PM_{2.5}$  to which they were exposed as modelled by the household  $PM_{2.5}$  mapping function for each location and year. Because indoor  $PM_{2.5}$  concentrations are consistently higher than those measured by personal monitors, we developed a ratio to scale the indoor exposure to female exposure. We then used female-to-male and female-to-child ratios that we developed to estimate exposure for males and children, respectively (appendix 1 pp 10–14). RR values for the corresponding overall exposure were

derived from the exposure–response curve (appendix 1 pp 14–44) and used to calculate PAFs (appendix 1 pp 45–46). We did not assume that all HAP exposure occurred at high concentrations of  $PM_{2.5}$ , which would have resulted in negligible changes in RR per unit change of  $PM_{2.5}$  concentration for much of the domain of exposure; rather, we estimated the PAF for HAP on the basis of the proportion of the overall (HAP plus ambient air pollution) exposure due to HAP. This approach also avoided the potential for double counting of the ambient contribution to HAP that is estimated from the household  $PM_{2.5}$  mapping function. We estimated the attributable burden for each cause as the product of the total burden for that cause and corresponding PAF for each GBD location, year, age group, and sex. To account for uncertainties in our PAF modelling, we produced 500 simulations of all estimates and intermediate steps (appendix 2 p 7). The PAF reported is the mean of these simulations, and the uncertainty interval was calculated as the 2.5th and 97.5th percentile of the simulations. Finally, we conducted decomposition analysis<sup>11</sup> to understand the causes of the observed changes in burden.

**Role of the funding source**

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all



**Figure 1: Percentage of global population exposed to HAP from solid cooking fuels, 2021**  
 HAP=household air pollution.

the data in the study and had final responsibility for the decision to submit for publication.

## Results

### Exposure to solid fuels

In 2021, 2.67 billion (95% uncertainty interval [UI] 2.63–2.71) people were exposed to HAP from solid cooking fuels, 33.8% (95% UI 33.2–34.3) of the global population (figure 1). Although this percentage is smaller than the 56.7% (56.4–57.1) of the global population exposed to solid fuels in 1990, the total number exposed has declined by only 0.35 billion (11.6%) since then (from 3.02 billion [95% UI 3.01–3.05]). Percentages of populations exposed to solid cooking fuels in 2021 remained highest in sub-Saharan Africa (78.8%, 95% UI 77.6–80.1), south Asia (53.2%, 51.9–54.5), and southeast Asia, east Asia, and Oceania (29.4%, 28.1–30.7; figure 1). In the Americas, Haiti was an outlier, with 91.3% (89.0–93.3) of individuals exposed to HAP in 2021 (figure 1).

Fuel-specific modelling showed that wood was the most prevalent solid fuel worldwide, with 24.0% (23.5–24.5) of the global population exposed in 2021 (table). Coal was the second most prevalent fuel globally, with 5.4% (5.2–5.6) of the population exposed in 2021.

Temporal trends showing the percentage of population exposed to each fuel type by super-region<sup>22</sup> and year are shown in figure 2. Although exposure to wood remained widespread globally, exposure to coal was lower, being mainly limited to sub-Saharan Africa and southeast Asia, east Asia, and Oceania (geographical distributions are in appendix 1 p 47).

Using crop residues as cooking fuel resulted in exposure to the highest concentrations of  $PM_{2.5}$ , with an exponentiated  $\beta$  value of 23.3 compared with the reference (ie, clean fuel), corresponding to exposure to concentrations of  $PM_{2.5}$  23.3 times higher than those produced by clean fuel, followed by dung ( $\beta$  value 10.5), wood ( $\beta$  value 7.3), and coal ( $\beta$  value 5.3; appendix 1 p 12). The mean concentration of  $PM_{2.5}$  from HAP globally was 213.6  $\mu\text{g}/\text{m}^3$  (median 198.0  $\mu\text{g}/\text{m}^3$ ) in 1990 and 84.2  $\mu\text{g}/\text{m}^3$  (median 61.5  $\mu\text{g}/\text{m}^3$ ) in 2021. The maximum concentration of  $PM_{2.5}$  from any solid cooking fuel to which individuals were exposed between 1990 and 2021 was 1660  $\mu\text{g}/\text{m}^3$ .

### RRs from HAP exposure

Figure 3 shows splines characterising the exposure–response relationships between  $PM_{2.5}$  exposure and COPD, ischaemic heart disease, lower respiratory





Figure 2: Percentages of population exposed to solid fuel types, globally and by super-region, 1990–2021





**Figure 3: Exposure–response relationships for  $PM_{2.5}$  concentration and outcomes except for cataract**

Risk curves were calculated on the basis of epidemiological data characterising exposure to ambient  $PM_{2.5}$  air pollution, and household air pollution. Shading indicates 95% uncertainty interval; x-axis truncated at  $600 \mu\text{g}/\text{m}^3$  to provide more detail at lower concentrations. COPD=chronic obstructive pulmonary disease.

infections, cancers of the trachea, bronchus, and lung, stroke, and type 2 diabetes. Across the full domain of exposure, COPD showed the highest RR from HAP exposure and type 2 diabetes the lowest. For cataract, with the burden of proof risk function approach and solid fuel use treated as a dichotomous risk factor (appendix 1 p 39), RR was 2.52 (95% CI 1.36–4.50). Curves for low birthweight and short gestation are in appendix 1 (pp 40–41).

### Burden

In 2021, 111 million (95% UI 75.1–164) disability-adjusted life-years (DALYs), 3.9% (95% UI 2.6–5.7) of the global burden from all risks, and 3.11 million (95% UI 1.90–5.19) deaths were attributable to HAP globally. HAP-attributable burden remained highest in sub-Saharan Africa and south Asia, with 4044.1 (3103.4–5219.7) and 3213.5 (2165.4–4409.4) age-standardised DALYs per 100000 population, respectively (table, figure 4). Comparing sexes, females were exposed to higher  $PM_{2.5}$  concentrations than males (appendix 1 p 12),<sup>16</sup> but, due to the higher burden of cardiovascular disease in males, HAP-attributable burden was slightly higher for males (appendix 1 p 48). All-cause PAFs for HAP have declined considerably since 1990 (appendix 1 pp 49–51), but HAP remains an important risk factor, with almost 30% of cataract burden and almost 20% of COPD burden due to HAP in 2021. Moreover, more than 0.5 million deaths in children younger than 5 years could be attributed to HAP, highlighting that almost 11% of under-5 mortality is due to HAP.

Although the percentage of global HAP-attributable DALYs has declined continuously over the past

three decades, our study found that one in 26 DALYs worldwide can still be attributed to HAP. Among Level 4 risks, HAP declined from being the second most important risk for deaths in 1990 to being the eighth most important in 2021, and from being the third most important risk for DALYs in 1990 to being the eighth most important in 2021.<sup>11</sup> The Socio-demographic Index<sup>23</sup> (SDI; a measure that captures income per capita, education, and fertility) is predictive of the proportion of a population experiencing burden due to HAP, but there is variation within regions and SDI levels (appendix 1 p 52). Notably, many locations in southeast Asia, east Asia, and Oceania shared a similar level of burden to locations in sub-Saharan Africa, despite having higher SDI values. Additionally, countries such as Somalia and Niger had low PAFs for HAP despite their low SDI values. In contrast, PAFs for HAP were high in nations such as the Solomon Islands and Vanuatu, even though their SDIs fall within the medium range.


In 2021, 518.1 (95% UI 410.1–641.7) age-standardised DALYs per 100000, about one-third of the DALYs attributable to HAP, were mediated through low birthweight and short gestation (figure 5). These mediated causes included otitis, meningitis, encephalitis, sudden infant death syndrome, upper respiratory infections, diarrhoeal diseases, neonatal disorders, and some of the lower respiratory burden. Of the Level 3 causes, ischaemic heart disease (18.2 million [95% UI 10.3–31.4]), stroke (18.2 million [10.7–30.9]), and COPD (15.6 million [9.6–25.3]) accounted for most of the direct HAP-attributable DALYs, whereas neonatal disorders were the leading cause of mediated HAP-attributable DALYs (32.1 million [25.4–39.7]). Years of life lost (YLLs) due to premature mortality and years lived with disability (YLDs), which are summed to get DALYs, showed very different patterns. Whereas YLDs consisted almost entirely of direct burden, particularly COPD (2.37 million [1.38–3.98]), cataract (1.96 million [0.612–4.03]), and type 2 diabetes (1.71 million [0.711–3.44]), the distribution of YLLs was almost evenly split between direct and mediated causes, with neonatal disorders making up the largest proportion of burden (32.0 million [25.4–39.7]). Deaths attributable to HAP were heavily dominated by direct causes, particularly ischaemic heart disease (0.763 million [0.413–1.37]), stroke (0.758 [0.433–1.31]), and COPD (0.694 million [0.412–1.18]), although neonatal disorders also contributed a notable proportion of mortality (0.356 million [0.282–0.441]).

We conducted decomposition analysis to examine the changes in burden attributable to HAP from 1990 to 2021 (figure 6). This method decomposes changes to attributable burden into percentages due to population growth, population ageing, change in exposure to the risk factor, and risk-deleted burden (ie, all other changes in burden not explained by the first three categories). The result shows an overall decline in burden globally that reflects the decline in exposure for all super-regions. However, this decline has been counteracted by population growth in

	DALYs (95% UI), 2021													
	Percentage of population exposed (95% UI)					DALYs (95% UI), 2021								
	Solid cooking fuels, 1990	Solid cooking fuels, 2010	Solid cooking fuels, 2021	Coal and charcoal, 1990	Coal and charcoal, 2021	Crop residues, 1990	Crop residues, 2021	Dung, 1990	Dung, 2021	Wood, 1990	Wood, 2021	Number (in millions), all ages	DALYs (%), all ages	DALYs, rate (per 100000), age standardised
Global	56.7% (56.4-57.1)	44.3% (44.1-44.6)	33.8% (33.3-34.3)	10.2% (9.9-10.5)	5.4% (5.2-5.6)	2.4% (2.0-2.7)	2.6% (2.3-2.9)	1.9% (1.7-2.2)	1.7% (1.5-2)	42.3% (41.7-42.8)	24.0% (23.5-24.5)	111.0 (75.1-164.0)	3.9% (2.6-5.7)	1500.3 (1028.4-2195.6)
Central Europe, eastern Europe, and central Asia	19.4% (17.9-21)	11.0% (10.2-11.8)	7.5% (6.8-8.3)	1.8% (1.4-2.2)	0.628% (0.508-0.796)	0.341% (0.233-0.471)	0.213% (0.142-0.304)	0.533% (0.404-0.683)	0.324% (0.262-0.398)	16.8% (15.4-18.3)	6.3% (5.7-7.0)	0.835 (0.277-2.77)	0.441% (0.149-1.472)	173.8 (68.4-509.8)
High-income countries	1.8% (1.6-1.9)	0.595% (0.533-0.662)	0.397% (0.340-0.460)	0.141% (0.108-0.183)	0.042% (0.034-0.053)	0.308% (0.285-0.332)	0.088% (0.059-0.127)	0.003% (0.002-0.005)	0.003% (0.002-0.006)	1.3% (1.2-1.4)	0.264% (0.222-0.313)	0.0194 (0.000377-0.139)	0.005% (0.0-0.041)	1.0 (0-7.8)
Latin America and Caribbean	32.5% (30.9-34.0)	18.2% (17.1-19.6)	11.7% (10.7-12.9)	1.6% (1.4-1.8)	1.6% (1.4-1.8)	0.012% (0.007-0.020)	0.015% (0.010-0.021)	0.121% (0.084-0.169)	0.059% (0.044-0.081)	30.8% (29.2-32.3)	10.1% (9.1-11.2)	1.79 (0.977-3.52)	0.853% (0.467-1.7)	316.7 (176.4-607.4)
North Africa and Middle East	18.4% (17.1-19.9)	11.0% (10.6-11.3)	6.6% (6.0-7.1)	2.0% (1.6-2.5)	0.793% (0.607-1.0)	1.5% (1.1-2.0)	1.2% (0.9-1.5)	1.4% (1.0-1.9)	0.910% (0.712-1.1)	13.4% (12.3-14.8)	3.7% (3.2-4.1)	2.5 (1.8-3.3)	1.3% (0.914-1.7)	471.9 (337.7-637.8)
South Asia	84.7% (84.0-85.4)	70.6% (70.1-71.0)	53.2% (51.9-54.5)	2.4% (2.1-2.6)	1.1% (0.947-1.2)	8.6% (7.1-10.2)	9.1% (7.8-10.3)	8.6% (7.5-9.8)	6.9% (5.7-8.1)	65.2% (63.4-67.0)	36.2% (34.5-37.8)	47.2% (31.9-64.9)	6.8% (4.6-9.3)	3213.5 (2165.4-4409.5)
Southeast Asia, east Asia, and Oceania	80.8% (80.0-81.5)	51.5% (50.7-52.4)	29.4% (28.1-30.7)	25.2% (24.4-26.1)	10.0% (9.2-10.7)	1.0% (0.814-1.2)	0.147% (0.111-0.198)	0.002% (0.002-0.002)	0.001% (0.001-0.002)	54.5% (53.5-55.6)	19.3% (18.2-20.4)	20.0 (7.93-45.6)	3.0% (1.2-7.0)	836.5 (350.4-1841.5)
Sub-Saharan Africa	90.8% (90.1-91.4)	84.3% (83.9-84.7)	78.8% (77.6-80.1)	13.8% (12.9-14.9)	15.2% (14.5-15.9)	1.0% (0.706-1.4)	2.1% (1.8-2.5)	0.326% (0.268-0.392)	0.287% (0.227-0.359)	75.7% (74.5-76.8)	61.3% (59.9-2.6)	39.2 (29.0-51.0)	6.7% (5.0-8.6)	4044.1 (3103.5-5219.5)


Some percentages and 95% UI values given to three decimal places for greater accuracy. Only solid cooking fuel is reported for 2010. DALYs=disability-adjusted life-years. HAP=household air pollution. UI=uncertainty interval.

**Table: Percentage of population exposed to HAP in 1990, 2010, and 2021 and number, percentage, and rate of DALYs attributable to HAP in 2021, globally and by super-region**



**Figure 4:** Population attributable fraction of DALYs attributable to HAP (A) and age-standardised rate per 100000 population of DALYs attributable to HAP (B), by location, 2021  
 DALY=disability-adjusted life-years. HAP=household air pollution.






**Figure 5: Composition of global, HAP-attributable Level 3 DALYs (A), deaths (B), YLLs (C), and YLDs (D) by disease, 2021**  
 Non-communicable diseases are in blue; communicable, maternal, neonatal, and nutritional diseases are in red. COPD=chronic obstructive pulmonary disease. DALYs=disability-adjusted life-years. HAP=household air pollution. YLD=years lived with disability. YLL=years of life lost.

almost every super-region, especially sub-Saharan Africa, whereas population ageing has caused a reduction of HAP-attributable DALYs in some super-regions.

## Discussion

Our study presents timely and comprehensive estimates of exposure to  $PM_{2.5}$  pollution from HAP and of the HAP-attributable disease burden in 2021. The findings show that, first, exposure to HAP from solid fuels remained very high in sub-Saharan Africa and south Asia and, although the percentage of the global population exposed to HAP declined between 1990 and 2021, the total number of individuals exposed to HAP in 2021 remained similar to the number estimated for 1990. Second, a substantial share of the burden of disease—almost a third of DALYs—attributable to HAP was mediated through adverse reproductive outcomes, with most of this burden comprising YLLs caused by neonatal disorders. Third, even accounting for the mediated burden, overall burden from HAP is higher than previously estimated, and HAP remained a leading risk factor globally for morbidity and mortality in 2021, with 111 million (95% UI 75.1–164) DALYs and 3.1 million (1.9–5.2) deaths attributable to HAP in 2021.

Estimates of HAP exposure from solid fuels in this study were similar to previous estimates. In 2014, Bonjour and colleagues,<sup>2</sup> detailing the exposure estimation methods used for GBD 2010, reported that 2.8 billion people were exposed to HAP (ie, 41% of the global population). Our analysis for GBD 2021 estimated that 3.08 billion (96% UI 3.06–3.10) people were exposed to HAP from solid fuels in 2010 (ie, 44.3% of the global population). The difference is probably the result of differing sources and modelling approaches. In GBD 2010, a single exposure indicator was used for all individuals exposed to HAP, regardless of year and location, whereas we used a spatially and temporally



**Figure 6: Decomposition of all-cause HAP-attributable DALYs into percent change driven by population growth, population ageing, exposure, or risk-deleted DALY rate, globally and by super-region, 1990–2021**  
 DALYs=disability-adjusted life-years. HAP=household air pollution.

variable model that accounted for the type of fuel used. Additionally, Bonjour and colleagues<sup>2</sup> used 586 country-year datapoints from 155 countries, whereas we used 1173 country-year sources from 161 countries (appendix 1 pp 5–7, appendix 2 p 1). A more recent study by Stoner and colleagues<sup>7</sup> used a multivariate hierarchical approach to estimate the proportion of individuals exposed to polluting fuels from 1990 to 2030; these estimates comprise those reported and used by WHO<sup>1</sup> for their estimates of disease burden.<sup>24</sup> Stoner and colleagues' model used six fuel types: kerosene, wood, crop waste,

dung (squeezed to biomass), charcoal, and coal.<sup>7</sup> As in our study, their model constrained the proportions of fuel types present so that the total never exceeded 100%. Stoner and colleagues used data already tabulated in the WHO Household Energy Database, whereas we extracted the microdata ourselves when possible and used tabulated estimates from the WHO Household Energy Database only when microdata were not available. This approach allowed us to ensure consistency in mapping survey responses to fuel categories and to extract household size, which we used to adjust results reported at the household level to a population estimate. Stoner and colleagues' estimate of the number of people exposed to polluting fuels in 2010 (3.0 billion [95% UI 2.7–3.3]) was very similar to our own, but their estimate for 2020 (2.8 billion [2.3–3.3] people exposed, 36% [95% UI 30–43] of the global population) was higher than our estimate for the same year (2.71 billion [95% UI 2.68–2.75] people exposed, 34.7% [95% UI 34.3–35.2] of the global population). This variation is probably due to modelling differences and the fact that we adjusted for household size. Nonetheless, our estimates for 2020 were within the uncertainty reported by Stoner and colleagues, and our estimates would probably have increased if we had included kerosene in our model.

Our study found a consistently higher HAP-attributable burden than previously estimated. In the finalised results for GBD 2010, we reported 108 million (95% UI 84.9–133) DALYs attributable to HAP in 2010,<sup>25</sup> whereas, here, we estimated 161 million (115–211) DALYs for the same year. Our estimate of deaths attributable to HAP in 2010 (4.23 million [2.94–5.77]) in this study is also higher than the number estimated in GBD 2010 (3.48 million [2.64–4.39]).<sup>25</sup> These differences can probably be explained by our updated methodology, which included differing estimation methods for  $PM_{2.5}$  concentrations and risk. Since 2010, we have added cataract and type 2 diabetes,<sup>17</sup> and, most notably, we have attributed a portion of the burden from causes mediated via adverse reproductive outcomes to HAP,<sup>18</sup> which has led to a substantial increase in our overall estimated burden. In addition, many of the epidemiological studies used in GBD 2010, especially for non-communicable diseases, were conducted in low-pollution settings and, thus, underestimated risk when informing burden for individuals exposed to high concentrations of pollution. Finally, we estimated both proportion of population exposed to HAP and  $PM_{2.5}$  concentrations from solid fuels on the basis of the specific type of fuel used. Our analysis showed that the burning of crop residue produces higher quantities of particulate matter than does burning of dung, wood, or coal (listed in descending order of the quantities of emissions produced).

A study of HAP-attributable burden by Lee and colleagues<sup>8</sup> used a random effects model to produce pooled estimates of RR for cardiorespiratory, maternal, and paediatric outcomes. Lee and colleagues estimated

that HAP contributed to 1.8 million (95% CI 1.1–2.7) deaths and 60.9 million (34.6–93.3) DALYs in 2017, whereas, here, we estimated 3.25 million (95% UI 2.01–5.14) deaths and 123 million (82.1–176) DALYs attributable to HAP in the same year. Our study estimated the actual concentrations of  $PM_{2.5}$  to which individuals were exposed and assigned excess risk, whereas Lee and colleagues used a binary exposure model (ie, exposed *vs* unexposed individuals) to estimate RR. The binary exposure model included kerosene and  $NO_2$  from cooking fuels, whereas our risk definition included  $PM_{2.5}$  pollution from solid fuels only. As our study reflects geographical, temporal, and fuel-type variations, we believe that our estimates reflect exposure more accurately. Lee and colleagues used the exposure estimates reported by Bonjour and colleagues;<sup>2</sup> thus, most of the additional burden in our analysis was probably driven by our higher estimates of exposure compared with the GBD 2010 study. Other differences were that, unlike Lee and colleagues, we included type 2 diabetes and causes mediated via adverse reproductive outcomes (which, together, constitute almost half of the estimated DALYs) and we did not model pulmonary tuberculosis or asthma. The inclusion of asthma symptoms and exacerbations is inconsistent with current risk factor analysis for air pollution in GBD, which focuses on incident disease, given the challenges in estimating short-term (ie, daily) variation in global exposure. Future iterations of the GBD study might include tuberculosis if forthcoming burden of proof analyses meet GBD inclusion criteria.<sup>19</sup> Finally, we derived our RR estimates from longitudinal and case-control studies and excluded cross-sectional data (appendix 1 p 39), whereas Lee and colleagues used cross-sectional data in their estimation.

By modelling fuel-specific exposure to HAP and then mapping location-specific and year-specific exposure to real-world measurements of  $PM_{2.5}$  by age and sex, our exposure estimation overcomes some of the limitations of previous analyses. Although our previous estimates relied on an integrated exposure–response curve,<sup>10</sup> our latest estimates used RR curves produced with the burden of proof risk function approach,<sup>19</sup> allowing us to adjust the RR input data for confounding covariates and interstudy heterogeneity. The availability of additional epidemiological studies specifically linking ambient pollution and HAP to morbidity and mortality also allowed us to discontinue the inclusion of active and passive smoking data in our RR estimates. The inclusion of HAP-specific studies, particularly those for cardiovascular disease, stroke, and type 2 diabetes, is a major improvement over our 2010 estimates. The decomposition analysis revealed that population growth has partly counteracted the effect of reduced exposure to HAP in almost every super-region and at the global level.

As is the case with other estimates of global risk burden, our estimates have several limitations. First, our

risk definition—exposure to PM<sub>2.5</sub> pollution from solid fuels—did not allow us to capture other sources of HAP that have established epidemiological effects, such as kerosene,<sup>26,27</sup> or co-pollutants, such as carbon monoxide.<sup>14,28</sup> WHO estimated that in 2021, 63·1 million (95% UI 37·8–120·4) people cooked primarily with kerosene.<sup>29</sup> Pollution produced by heating and lighting fuels in the home was absent from our analysis, which probably resulted in overestimation of the specific effect of pollution from cooking fuels. We also encountered several challenges concerning the input data for our exposure models. As we extracted only data representing the primary fuel used in a home, and data on secondary fuel sources are sparse, we were unable to model usage of multiple fuel types (known as stove stacking), which remains highly prevalent among communities using solid fuels.<sup>30,31</sup> Exposure data were also very sparse for high-income countries, eastern Europe, and Oceania; thus, estimates for these locations relied heavily on data imputation through spatiotemporal Gaussian process regression inference and might have underestimated solid fuel usage for communities who remain without access to cleaner fuels in these locations. Our estimates did not include the effects of improved stoves, although the efficacy of such interventions seems low.<sup>32–35</sup> Similarly, we were unable to capture the effects of various grades of fuel within a category; instead, we assumed equitoxicity within each fuel type. Furthermore, the uncertainty of our RR model could not account for measurement bias, selection bias, or model mis-specification bias. Another important limitation is the fact that we did not adjust for confounding bias because our PAF estimation method assumed no unmeasured confounding. The wide UIs for our RR curves, particularly at high PM<sub>2.5</sub> concentrations, indicate the need for more high-quality studies. Dementia and tuberculosis were not included in our estimations, but these outcomes might be evaluated for inclusion in future GBD cycles.<sup>36,37</sup> Further discussion of the limitations of this study can be found in appendix 1 (pp 7, 10, 15). In summary, these omissions probably led to further underestimation of exposure, and, thus, burden in this study.

Our study confirms that, although progress has been made in reducing exposure to and burden attributable to HAP, the rapid transition of under-resourced communities to cleaner fuel sources is urgently needed, given the magnitude of current attributable disease burden. Reductions in exposure have been geographically variable, with the smallest declines in sub-Saharan Africa and south Asia. Previous studies have focused on household-level interventions, such as improved stoves, but the limited success of these programmes indicates the need for investments in community-level infrastructure. For example, the recent Household Air Pollution Intervention Network (HAPIN) trial<sup>38</sup>—the largest randomised controlled trial to study the switching of cooking fuels from biomass to liquefied petroleum

gas—indicated reductions in exposure to PM<sub>2.5</sub>, but the intervention did not lead to reduced incidence of childhood pneumonia or to increased birthweight.<sup>39,40</sup> Importantly, in the intervention group, the median PM<sub>2.5</sub> concentration was 24·2 µg/m<sup>3</sup>,<sup>39</sup> a concentration for which our analysis still indicated substantially increased risk of low birthweight (appendix 1 pp 40–44), a possible limitation acknowledged by the authors of the HAPIN trial.<sup>38</sup> Like the HAPIN trial, the Ghana Randomized Air Pollution and Health Study (GRAPHS) found that switching to liquefied petroleum gas produced no improvement in birthweight or the incidence of severe pneumonia.<sup>35</sup> The median PM<sub>2.5</sub> concentration to which the intervention group in GRAPHS was exposed was 45 µg/m<sup>3</sup>, probably due to pollution from neighbours' cooking. A systematic review by Puzzolo and colleagues<sup>41</sup> of respiratory outcomes, low birthweight, and short gestation found benefits to cooking with gas rather than solid fuels but also found that the use of gas instead of electric stoves increased the risk of pneumonia and COPD. Unlike our study, the meta-analysis by Puzzolo and colleagues included studies on heating fuel usage and cross-sectional studies; they note that cross-sectional studies provide a lower level of evidence but point to the paucity of randomised controlled trials available for inclusion. Due to the timing of the search, Puzzolo and colleagues did not include the HAPIN findings in their systematic review, but they acknowledge the study on birthweight<sup>40</sup> and reiterate a limitation reported by the HAPIN authors—ie, the fuel switch was not initiated until the second trimester of pregnancy, possibly limiting the efficacy of the intervention.<sup>39,40</sup> These contrasting findings highlight the complexity of the healthy household energy challenge and make clear that further, high-quality studies are needed, particularly studies that focus on community-level intervention.

The high level of paediatric burden estimated in this study is a major cause for concern. Apart from the approximately 66 million DALYs of HAP-attributable burden mediated via adverse reproductive outcomes in 2021, paediatric exposure to PM<sub>2.5</sub> has lifelong implications, including developmental disorders, IQ loss, and increased risk of chronic illness in adulthood.<sup>42</sup> Paediatric HAP exposure also impairs educational attainment and lifetime economic potential, compounding the disadvantages already faced by children from low-income backgrounds.<sup>43</sup> Although several measurement studies have indicated that females are exposed to higher concentrations of particulate matter from HAP than are males (appendix 1 pp 11–13), these differences are less pronounced than often believed. For our study, we calculated adjustment ratios on the basis of measurements in several studies, including those in the PURE-AIR study, which measured exposure to household and personal air pollution within 120 communities in eight countries.<sup>16</sup> The ratios we estimated were 0·64 (95% CI 0·52–0·79) for males and 0·85 (0·67–1·09) for children. The difference observed in

the PURE-AIR study was even less pronounced, with mean  $PM_{2.5}$  concentrations of  $67 \mu\text{g}/\text{m}^3$  (95% CI 62–72) for females and  $62 \mu\text{g}/\text{m}^3$  (58–67) for males. The overall HAP-attributable burden was higher for males than for females; this finding is the consequence of an overall higher disease burden in males than in females, especially for cardiometabolic disease. Policy makers should note that although cooking is often done by women,<sup>5</sup> leading to increased exposure to HAP, all members of exposed households are harmed by HAP. HAP also makes a large contribution to ambient air pollution,<sup>44,45</sup> and the elimination of HAP could help affected countries to meet their goals for ambient air quality.<sup>46,47</sup> For example, in India, 32% of the overall contribution of residential combustion to mortality was mediated via its effect on exposure to ambient  $PM_{2.5}$ .<sup>48</sup> In addition, the inefficient combustion of solid cooking fuels releases black carbon and  $CO_2$ ,<sup>16</sup> climate forcing agents that contribute to anthropogenic climate change.<sup>49</sup> Floess and colleagues<sup>50</sup> have shown that transitioning users of solid fuels to cleaner energy sources also has large co-benefits for reduction of greenhouse gas emissions. Thus, the positive effects of equitable and culturally sensitive programmes in transitioning communities from solid cooking fuels to cleaner fuels are numerous. Renewed international investment to provide under-resourced communities, particularly those in sub-Saharan Africa and south Asia, with cleaner fuels must be renewed and accelerated.

In conclusion, HAP remains a leading risk factor in many low-income and middle-income countries, exerting adverse effects on a variety of health outcomes, including cardiorespiratory and metabolic diseases, as well as on reproductive health outcomes. Our study underscores the serious health consequences linked to adverse reproductive health outcomes; these can be both fatal and long lasting and have not received adequate attention from researchers and policy makers. We hope our work further illuminates the fact that the HAP-derived burden remains a major problem for many low-income and middle-income countries. Despite steady improvements made over recent years, we argue that HAP mitigation needs to remain high up the global policy agenda and requires a multifaceted approach. Prioritising research and development for cleaner technologies, subsidising affordable clean energy appliances, and enforcing stringent emissions standards are crucial policy measures. Equally important are public awareness campaigns that educate people about the associated health risks of HAP, alongside enhanced monitoring systems that supply the data needed for evidence-based decision making. By incentivising private sector investment and ensuring cross-sector policy integration, we can establish comprehensive and effective strategies to substantially reduce HAP and its detrimental effects.<sup>1,51</sup>

#### GBD 2021 HAP Collaborators

Fiona B Bennitt, Sarah Wozniak, Kate Causey, Sandra Spearman, Chukwuma Okereke, Vanessa Garcia, Nadim Hashmeh,

Charlie Ashbaugh, Atef Abdelkader, Meriem Abdoun, Muhammed Jemal Abdurebi, Armita Abedi, Roberto Ariel Abeldaño Zuñiga, Richard Gyan Aboagye, Bilyaminu Abubakar, Ahmed Abu-Zaid, Mesafint Molla Adane, Oyelola A Adegbeye, Victor Adekanmbi, Abiola Victor Adepoju, Temitayo Esther Adeyeoluwa, Olorunsola Israel Adeyomoye, Rishan Adha, Muhammad Sohail Afzal, Saira Afzal, Feleke Doyore Agide, Aqeel Ahmad, Danish Ahmad, Muayyad M Ahmad, Sajjad Ahmad, Ali Ahmadi, Sepideh Ahmadi, Anisuddin Ahmed, Ayman Ahmed, Haroon Ahmed, Marjan Ajami, Rufus Olusola Akinyemi, Salah Al Awaidey, Hanadi Al Hamad, Muaaz M Alajlani, Mulubirhan Assefa Alemayohu, Adel Ali Saeed Al-Gheethi, Abid Ali, Waad Ali, Sheikh Mohammad Alif, Sami Almustanyir, Nelson Alvis-Guzman, Nelson J Alvis-Zakzuk, Hany Aly, Hubert Amu, Ganiyu Adeniyi Amusa, Tadele Fentabel Anagaw, Boluwatife Stephen Anuoluwa, Iyadunni Adesola Anuoluwa, Saeid Anvari, Ekenedilichukwu Emmanuel Anyabolo, Geminn Louis Carace Apostol, Aleksandr Y Aravkin, Demelash Arede, Brhane Berhe Aregawi, Olatunde Aremu, Akeza Awealom Asgedom, Mubarek Yesse Ashemo, Tahira Ashraf, Seyyed Shamsadin Athari, Sina Azadnajafabad, Ahmed Y Azzam, Giridhara Rathnaiah Babu, Saeed Bahramian, Kiran Bam, Maciej Banach, Biswajit Banik, Mehmet Firat Baran, Francesco Barone-Adesi, Sandra Barteit, Hameed Akande Bashiru, Pritish Baskaran, Mohammad-Mahdi Bastan, Sanjay Basu, Saurav Basu, Sefealem Assefa Belay, Melesse Belayneh, Apostolos Beloukas, Derrick A Bennett, Devidas S Bhagat, Dinesh Bhandari, Pankaj Bhardwaj, Sonu Bhaskar, Ajay Nagesh Bhat, Priyadarshini Bhattacharjee, Gurjit Kaur Bhatti, Manpreet S Singh Bhatti, Cem Bilgin, Mary Sefa Boampong, Sri Harsha Boppana, Samuel Adolf Bosoka, Sofiane Boudalia, Fan Cao, Rama Mohan Chandika, Gashaw Sisay Chanie, Vijay Kumar Chattu, Anis Ahmad Chaudhary, Akhilanand Chaurasia, Guangjin Chen, Yifan Chen, Ritesh Chimoriya, Bryan Chong, Devasahayam J Christopher, Isaac Sunday Chukwu, Aaron J Cohen, Natalia Cruz-Martins, Omid Dadras, Xiaochen Dai, Patience Unekwojo Daikwo, Samuel Demissie Darcho, Saswati Das, Juana Maria Delgado-Saborit, Belay Desye, Sagnik Dey, Meghnath Dhimal, Daniel Diaz, Thanh Chi Do, Ojas Prakashbhai Doshi, Abdel Rahman E'mar, Alireza Ebrahimi, Hisham Atan Edinur, Aziz Eftekharimehrabad, Temitope Cyrus Ekundayo, Ibrahim Farahat El Bayoumy, Syed Emdadul Haque, Theophilus I Emeto, Habtamu Demelash Enyew, Ayesha Fahim, Adekunle Gregory Fakunle, Sasan Faridi, Timur Fazlyov, Alireza Feizkhal, Florian Fischer, Morenike Oluwatoyin Folayan, Sridevi G, Muktar A Gadanya, Xiang Gao, Miglas Welay Gebregergis, Mesfin Gebrehiwot, Teferi Gebru Gebremeskel, Afsaneh Ghazemzadeh, Nermin Ghith, Mahaveer Golechha, Davide Golinelli, Shi-Yuan Guan, Zhifeng Guo, Bhawna Gupta, Lalit Gupta, Rabih Halwani, Ahmed I Hasaballah, Md Saquib Hasnain, Simon I Hay, Demisu Zenbaba Heyi, Kamal Hezam, Nguyen Quoc Hoan, Ramesh Holla, Hassan Hosseinzadeh, Chengxi Hu, Hong-Han Huynh, Bing-Fang Hwang, Segun Emmanuel Ibitoye, Oluwatope Olaniyi Idowu, Adalia Ikiroma, Mustapha Immurana, Arit Inok, Muhammad Iqhrammullah, Rakibul M Islam, Sheikh Mohammed Shariful Islam, Vinothini J, Ammar Abdulrahman Jairoun, Abhishek Jaiswal, Mihajlo Jakovljevic, Reza Jalilzadeh Yengejeh, Manthan Dilipkumar Janodia, Shubha Jayaram, Aleigh Tasew Jema, Ravi Prakash Jha, Jost B Jonas, Nitin Joseph, Vidya Kadashetti, Kehinde Kazeem Kanmodi, Sushil Kumar Kansal, Ibraheem M Karaye, Gbenga A Kayode, Himanshu Khajuria, Amirmohammad Khalaji, Vishnu Khanal, Khaled Khatib, Khalid A Kheirallah, Atulya Aman Khosla, Majid Khosravi, Shivakumar KM, Luke D Knibbs, Gerbrand Koren, Parvaiz A Koul, Kewal Krishan, Barthelemy Kuate Defo, Mohammed Kuddus, Mukhtar Kulimbet, Vishnutheertha Kulkarni, Ashish Kumar, Dewesh Kumar, Nithin Kumar, Om P Kurmi, Chandrakant Lahariya, Hanpeng Lai, Tuo Lan, Paolo Lauriola, Nhi Huu Hanh Le, Munjae Lee, Seung Won Lee, Stephen S Lim, Gang Liu, Shuke Liu, Wei Liu, José Francisco López-Gil, Jay B Lusk, Sandeep B Maharaj, Kashish Malhotra, Ahmad Azam Malik,



Iram Malik, Lesibana Anthony Malinga, Alexander G Mathioudakis, Rita Mattiello, Andrea Maugeri, Tesfahun Mekene Meto, Hadush Negash Meles, Ritesh G Menezes, Sultan Ayoub Meo, Seid Tiku Mereta, Tuomo J Meretoja, Tomislav Mestrovic, Laurette Mhlanga, Ted R Miller, Andreea Mirica, Erkin M Mirrakhimov, Moonis Mirza, Awoke Misganaw, Prasanna Mithra, Jama Mohamed, Nouh Saad Mohamed, Abdollah Mohammadian-Hafshejani, Mustapha Mohammed, Shafiu Mohammed, Ali H Mokdad, Shaher Momani, Himel Mondal, Lidia Morawska, Rohith Motappa, Sumaira Mubarik, Kavita Munjal, Yanjinlkhankham Munkhsaikhan, Christopher J L Murray, Woojae Myung, Sanjeev Nair, Vinay Nangia, Muhammad Naveed, Nawsherwan, Rawlance Ndeji, Dang Nguyen, Hien Quang Nguyen, Van Thanh Nguyen, Taxiarchis Konstantinos Nikolouzakakis, Vikram Niranjana, Efaq Ali Noman, Syed Toukir Ahmed Noor, Abbas Norouziyan Baghani, Jean Jacques Noubiap, Ogochukwu Janet Nzoputam, Bogdan Oancea, Ismail A Odetokun, Daniel Bogale Odo, Akinyemi O D Ofakunrin, Onome Bright Oghenetega, Osaretin Christabel Okonji, Andrew T Olagunju, Tosin Abiola Olasehinde, Isaac Iyinoluwa Olufadewa, Gideon Olamilekan Oluwatunase, Ahmed Omar Bali, Mohammad Mehdi Ommati, Abiodun Olusola Omotayo, Maureen Auma Ondayo, Adrian Otoi, Mayowa O Owolabi, Mahesh Padukudru P A, Jagadish Rao Padubidri, Ioannis Pantazopoulos, Shahina Pardhan, Pragyan Paramita Parija, Romil R Parikh, Eun-Keel Park, Ashwaghosha Parthasarathi, Jay Patel, Siddhartha Pati, Shrikant Pawar, Prince Peprah, Gavin Pereira, Arokiasamy Perianayagam, Hoang Tran Pham, Ramesh Poluru, Akram Pourshams, Jalandhar Pradhan, Elton Junio Sady Prates, Dimas Ria Angga Pribadi, Jagadeesh Puvvula, Ata Rafiee, Pankaja Raghav, Fakher Rahim, Mohammad Hifz Ur Rahman, Mosiur Rahman, Muhammad Aziz Rahman, Amir Masoud Rahmani, Mohammad Rahmanian, Sathish Rajaa, Rayan Rajabi, Prashant Rajput, Mahmoud Mohammed Ramadan, Jewel Rana, Kritika Rana, Chhabi Lal Ranabhat, Drona Prakash Rasali, Santosh Kumar Rauniyar, Salman Rawaf, Elrashdy M Moustafa Mohamed Redwan, Nazila Rezaei, Jefferson Antonio Buendia Rodriguez, Susanne Röhr, Gholamreza Roshandel, Himanshu Sekhar Rout, Priyanka Roy, Michele Russo, Cameron John Sabat, Basema Ahmad Saddik, Umar Saeed, Narjes Saheb Sharif-Askari, Amirhossein Sahebkar, Pragyan Monalisa Sahoo, Afeez Abolarinwa Salami, Dauda Salihu, Abdallah M Samy, Milena M Santric-Milicevic, Tanmay Sarkar, Maheswar Satpathy, Ganesh Kumar Saya, Md Abu Sayeed, Austin E Schumacher, Mihretu Tadesse Sergindo, Yashendra Sethi, Allen Seylani, Samiah Shahid, Sunder Sham, Muhammad Aaqib Shamim, Anas Shamsi, Aziz Sheikh, Pavanchand H Shetty, Aminu Shittu, Ivy Shiue, Emmanuel Edwar Siddig, Paramdeep Singh, Surjit Singh, Md Shahjahan Siraj, Jeffrey D Stanaway, Leo Stockfelt, Kurt Straif, Chandan Kumar Swain, Berwin Singh Swami Vetha, Seyyed Mohammad Tabatabaei, Mircea Tampa, Haos Tang, Manoj Tanwar, Elvis Enowbeyang Tarkang, Yibekal Manaye Tefera, Mohamad-Hani Temsah, Reem Mohamad Hani Temsah, Ramna Thakur, Friedrich Thienemann, Nigusie Selomon Tibebe, Krishna Tiwari, Marcos Roberto Tovani-Palone, Jaya Prasad Tripathy, Aristidis Tsatsakis, Munkhtuya Tumurkhuu, Aniefiok John Udoakang, Sana Ullah, Sanaz Vahdati, Siavash Vaziri, Madhur Verma, Simone Vidale, Simona Villani, Karn Vohra, Theo Vos, Gizachew Tadesse Wassie, Haftom Legesse Weldetinsaa, Adhena Ayalew Werkneh, Nuwan Darshana Wickramasinghe, Marcin W Wojewodzic, Tewodros Eshete Wonde, Felicia Wu, Zenghong Wu, Hong Xiao, Suowen Xu, Mukesh Kumar Yadav, Saba Yahoo Syed, Sanni Yaya, Arzu Yiğit, Vahit Yiğit, Dehui Yin, Dong Keon Yon, Naohiro Yonemoto, Chuanhua Yu, Leila Zaki, Mohammed G M Zeariya, Youjie Zeng, Chunxia Zhai, Haijun Zhang, Zhiqiang Zhang, Bin Zhu, Sa'ed H Zyoud, Samer H Zyoud, Michael Brauer, and Katrin Burkart. Please see appendix 3 (pp 3–15) for the affiliations for individual authors.

#### Contributors

Please see appendix 3 (pp 15–19) for more detailed information about individual author contributions to the research, divided into the following categories: managing the overall research enterprise; writing

the first draft of the manuscript; primary responsibility for applying analytical methods to produce estimates; primary responsibility for seeking, cataloguing, extracting, or cleaning data; designing or coding figures and tables; providing data or critical feedback on data sources; developing methods or computational machinery; providing critical feedback on methods or results; drafting the manuscript or revising it critically for important intellectual content; and managing the estimation or publications process.

#### Declaration of interests

S Afzal reports support for the present manuscript from King Edward Medical University for providing study material, research articles, valid data sources and authentic real time information. S Afzal reports payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from King Edward Medical University and collaborative partners including University of Johns Hopkins, University of California, University of Massachusetts, KEMCAANA, KEMCA UK; support for attending meetings and/or travel from King Edward Medical University; participation on a Data Safety Monitoring Board or Advisory Board with National Bioethics Committee Pakistan, King Edward Medical University Ethical Review Board, Ethical Review Board Fatima Jinnah Medical University and Sir Ganga Ram Hospital, and Member Technical Working Group on Infectious Diseases to formulate guidelines; leadership or fiduciary roles in board, society, committee or advocacy groups, paid or unpaid with Pakistan Association of Medical Editors, Faculty of Public Health Royal Colleges UK (FFPH) as Fellow, Society of Prevention, Advocacy And Research, King Edward Medical University (SPARK), and Pakistan Society of Infectious Diseases as a member; other support from King Edward Medical University as Dean of Public Health and Preventative Medicine and Director of Quality Enhancement Cell, Annals of King Edward Medical University as Chief Editor since 2014, Faculty of Public Health United Kingdom as Fellow, KEMCA-UK as Advisory Board Member and Chair Scientific Session, KEMCAANA as Chairperson International Scientific Conference, Research and Publications Higher Education Commission, HEC Pakistan as Member, Research and Journals Committee Paki-stan Medical and Dental Council, Pakistan as Member, National Bioethics Committee, Pakistan as Member, Punjab Corona Experts Advisory Group as Member, Technical Working Group on Infectious Diseases as Member, Dengue Experts Advisory Group as Member, Punjab Residency Program Research Committee as Chair; all outside the submitted work. S M Alif reports payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from Victoria University Online; support for attending meetings and/or travel from University of Melbourne; leadership or fiduciary roles in board, society, committee or advocacy groups, paid or unpaid with Thoracic Society of Australia and New Zealand; all outside the submitted work. S Barteit reports grants from Carl-Zeiss Foundation and German Research Foundation (DFG); stock or stock options in CHEERS company, a for-profit company focusing on climate change and health evaluation and response systems; all outside the submitted work. A Beloukas reports grants or sponsorships from Gilead and GSK/ViiV paid to the University of West Attica; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from Gilead and GSK/ViiV paid to the University of West Attica; support for attending meetings and/or travel from Gilead and GSK/ViiV; receipt of equipment, materials, drugs, medical writing, gifts or other services from Cepheid for FOC reagents provided for a research project; all outside the submitted work. S Bhaskar reports grant or contracts from Japan Society for the Promotion of Science (JSPS), Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) for a Grant-in-Aid for Scientific Research (KAKENHI) (23KF0126) and from JSPS and the Australian Academy of Science for the JSPS International Fellowship (P23712); leadership or fiduciary roles in board, society, committee or advocacy groups, paid or unpaid with Rotary District 9675, Sydney, Australia as District Chair, Diversity, Equity & Inclusion, Global Health & Migration Hub Community, Global Health Hub Germany, Berlin, Germany as Chair, Founding Member and Manager, PLOS One, BMC Neurology, Frontiers in Neurology, Frontiers in Stroke, Frontiers in Public Health, Journal of Aging Research & BMC Medical Research Methodology as Editorial Board Member, College of Reviewers, Canadian Institutes of

See Online for appendix 3

Health Research (CIHR), Government of Canada as Member, World Headache Society, Bengaluru, India as Director of Research; Cariplo Foundation, Milan, Italy as Expert Adviser/Reviewer, National Cerebral and Cardiovascular Center, Department of Neurology, Division of Cerebrovascular Medicine and Neurology, Suita, Osaka, Japan as Visiting Director, and Cardiff University Biobank, Cardiff, UK as Member, Scientific Review Committee; all outside the submitted work. T C Ekundayo reports grants or contracts from University of South Africa via a PDF research fellowship payment, outside the submitted work. N Ghith reports a grant from Novo Nordisk Foundation (NNF16OC0021856) via a salary during her employment at the Technical University of Denmark between 2019–2022; support for attending meetings and/or travel from Danish Data Science Institute at the Technical University of Denmark in 2023; all outside the submitted work. C Hu reports support for the present manuscript from the National Social Science Fund of China (grant number: 24CSH106). R M Islam reports other support from International Menopause Society (IMS) for serving as an Associate Editor of *Climacteric*, the official journal of the International Menopause Society (IMS), and as a member of the IMS Publication Steering Committee, outside the submitted work. K Krishan reports non-financial support from the UGC Centre of Advanced Study, CAS II, awarded to the Department of Anthropology, Panjab University, Chandigarh, India, outside the submitted work. M Lee reports support for the present manuscript from the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2023S1A3A2A05095298). S A Meo reports grants or contracts from Researchers Supporting Project, King Saud University, Riyadh, Saudi Arabia (RSP-2025 R47), outside the submitted work. B Oancea reports grants or contracts from the MRID (project PNR-18 no 842027778, contract no 760096), outside the submitted work. F Thienemann reports grants or contracts from Gates MRI, EDCTP, CRDF Global for studies at the University of Cape Town, EDCTP2 programme supported by the European Union (grant number RIA2017T-2004-StatintB), CRDF Global (G-202303-69943); outside the submitted work.

#### Data sharing

The data used in these analyses can be downloaded from the Global Health Data Exchange GBD 2021 website at <http://ghdx.healthdata.org/gbd-2021/sources>.

#### Acknowledgments

This research was funded by the Bill & Melinda Gates Foundation.

Editorial note: The Lancet Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

#### References

- International Energy Agency, International Renewable Energy Agency, UN Statistics Division, World Bank, WHO. Tracking SDG 7: the energy progress report. World Bank, 2024.
- Bonjour S, Adair-Rohani H, Wolf J, et al. Solid fuel use for household cooking: country and regional estimates for 1980–2010. *Env Health Perspect* 2013; **121**: 784–90.
- Smith KR, Bruce N, Balakrishnan K, et al. Millions dead: how do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution. *Ann Rev Public Health* 2014; **35**: 185–206.
- WHO. Guidelines for indoor air quality: household fuel combustion. Jan 21, 2014. <https://www.who.int/publications/i/item/9789241548885> (accessed Oct 31, 2024).
- WHO. Burning opportunity: clean household energy for health, sustainable development, and wellbeing of women and children. March 31, 2016. <https://www.who.int/publications/i/item/9789241565233> (accessed Oct 31, 2024).
- UN. Sustainable Development Goals. 2016. <https://www.un.org/sustainabledevelopment/> (accessed Oct 19, 2024).
- Stoner O, Lewis J, Martínez IL, Gumy S, Economou T, Adair-Rohani H. Household cooking fuel estimates at global and country level for 1990 to 2030. *Nat Commun* 2021; **12**: 5793.
- Lee KK, Bing R, Kiang J, et al. Adverse health effects associated with household air pollution: a systematic review, meta-analysis, and burden estimation study. *Lancet Glob Health* 2020; **8**: e1427–34.
- Shupler M, Godwin W, Frostad J, Gustafson P, Arku RE, Brauer M. Global estimation of exposure to fine particulate matter (PM<sub>2.5</sub>) from household air pollution. *Environ Int* 2018; **120**: 354–63.
- Stanaway JD, Afshin A, Gakidou E, et al. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018; **392**: 1923–94.
- Brauer M, Roth GA, Aravkin AY, et al. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet* 2024; **403**: 2162–203.
- Das Gupta P. Standardization and decomposition of rates: a user's manual. US Dept of Commerce, Economics and Statistics Administration, Bureau of the Census, 1993.
- WHO. Household energy database. <https://www.who.int/data/gho/data/themes/air-pollution/who-household-energy-db> (accessed July 5, 2019).
- Shupler M, Balakrishnan K, Ghosh S, et al. Global household air pollution database: kitchen concentrations and personal exposures of particulate matter and carbon monoxide. *Data Brief* 2018; **21**: 1292–95.
- WHO. Global database of household air pollution measurements. <https://www.who.int/data/gho/data/themes/air-pollution/hap-measurement-db> (accessed Nov 12, 2019).
- Shupler M, Hystad P, Birch A, et al. Household and personal air pollution exposure measurements from 120 communities in eight countries: results from the PURE-AIR study. *Lancet Planet Health* 2020; **4**: e451–62.
- Burkart K, Causey K, Cohen AJ, et al. Estimates, trends, and drivers of the global burden of type 2 diabetes attributable to PM<sub>2.5</sub> air pollution, 1990–2019: an analysis of data from the Global Burden of Disease Study 2019. *Lancet Planet Health* 2022; **6**: e586–600.
- Ghosh R, Causey K, Burkart K, Wozniak S, Cohen A, Brauer M. Ambient and household PM<sub>2.5</sub> pollution and adverse perinatal outcomes: a meta-regression and analysis of attributable global burden for 204 countries and territories. *PLoS Med* 2021; **18**: e1003718.
- Zheng P, Afshin A, Biryukov S, et al. The Burden of Proof studies: assessing the evidence of risk. *Nat Med* 2022; **28**: 2038–44.
- Zheng P, Barber R, Sorensen RJD, Murray CJL, Aravkin AY. Trimmed constrained mixed effects models: formulations and algorithms. *J Comput Graph Stat* 2021; **30**: 544–56.
- Murray CJL, Aravkin AY, Zheng P, et al. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; **396**: 1223–49.
- Institute for Health Metrics and Evaluation. Global Burden of Disease (GBD): which locations are studied in the GBD? <https://www.healthdata.org/research-analysis/about-gbd> (accessed Oct 24, 2024).
- Wang H, Abbas KM, Abbasifard M, et al. Global age-sex-specific fertility, mortality, healthy life expectancy (HALE), and population estimates in 204 countries and territories, 1950–2019: a comprehensive demographic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; **396**: 1160–203.
- WHO. Household air pollution. Oct 16, 2024. <https://www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health> (accessed Oct 19, 2024).
- Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2224–60.
- Lam NL, Smith KR, Gauthier A, Bates MN. Kerosene: a review of household uses and their hazards in low- and middle-income countries. *J Toxicol Environ Health B Crit Rev* 2012; **15**: 396–432.
- Arku RE, Brauer M, Duong M, et al. Adverse health impacts of cooking with kerosene: a multi-country analysis within the Prospective Urban and Rural Epidemiology Study. *Environ Res* 2020; **188**: 109851.
- Powers CI, Li L, Ezzati M, Butler JP, Zigler CM, Spengler JD. Chronic household air pollution and exposure patterns among Himalayan nomads. *J Expo Sci Environ Epidemiol* 2024; **34**: 973–80.

- 29 WHO. Population with primary reliance on fuels and technologies for cooking, by fuel type (in millions). July 1, 2024. <https://www.who.int/data/gho/data/indicators/indicator-details/GHO/population-with-primary-reliance-on-fuels-and-technologies-for-cooking-by-fuel-type> (accessed Nov 8, 2024).
- 30 Shankar AV, Quinn A, Dickinson KL, et al. Everybody stacks: lessons from household energy case studies to inform design principles for clean energy transitions. *Energy Policy* 2020; **141**: 111468.
- 31 Jewitt S, Atagher P, Clifford M. “We cannot stop cooking”: stove stacking, seasonality and the risky practices of household cookstove transitions in Nigeria. *Energy Res Soc Sci* 2020; **61**: 101340.
- 32 Pope D, Bruce N, Dherani M, Jagoe K, Rehfuess E. Real-life effectiveness of ‘improved’ stoves and clean fuels in reducing PM<sub>2.5</sub> and CO: systematic review and meta-analysis. *Environ Int* 2017; **101**: 7–18.
- 33 Quansah R, Semple S, Ochieng CA, et al. Effectiveness of interventions to reduce household air pollution and/or improve health in homes using solid fuel in low-and-middle income countries: a systematic review and meta-analysis. *Environ Int* 2017; **103**: 73–90.
- 34 Mortimer K, Ndamala CB, Naunje AW, et al. A cleaner burning biomass-fueled cookstove intervention to prevent pneumonia in children under 5 years old in rural Malawi (the Cooking and Pneumonia Study): a cluster randomised controlled trial. *Lancet* 2017; **389**: 167–75.
- 35 Jack DW, Ae-Ngibise KA, Gould CF, et al. A cluster randomised trial of cookstove interventions to improve infant health in Ghana. *BMJ Glob Health* 2021; **6**: e005599.
- 36 Pu F, Hu Y, Li C, et al. Association of solid fuel use with a risk score capturing dementia risk among middle-aged and older adults: a prospective cohort study. *Environ Res* 2023; **218**: 115022.
- 37 Hystad P, Larkin A, Rangarajan S, et al. Associations of outdoor fine particulate air pollution and cardiovascular disease in 157436 individuals from 21 high-income, middle-income, and low-income countries (PURE): a prospective cohort study. *Lancet Planet Health* 2020; **4**: e235–45.
- 38 Johnson M, Pillarisetti A, Piedrahita R, et al. Exposure contrasts of pregnant women during the Household Air Pollution Intervention Network randomized controlled trial. *Environ Health Perspect* 2022; **130**: 097005.
- 39 McCollum ED, McCracken JP, Kirby MA, et al. Liquefied petroleum gas or biomass cooking and severe infant pneumonia. *N Engl J Med* 2024; **390**: 32–43.
- 40 Clasen TF, Chang HH, Thompson LM, et al. Liquefied petroleum gas or biomass for cooking and effects on birth weight. *N Engl J Med* 2022; **387**: 1735–46.
- 41 Puzzolo E, Fleeman N, Lorenzetti F, et al. Estimated health effects from domestic use of gaseous fuels for cooking and heating in high-income, middle-income, and low-income countries: a systematic review and meta-analyses. *Lancet Respir Med* 2024; **12**: 281–93.
- 42 Brumberg HL, Karr CJ, Bole A, et al. Ambient air pollution: health hazards to children. *Pediatrics* 2021; **147**: e2021051484.
- 43 Case A, Fertig A, Paxson C. The lasting impact of childhood health and circumstance. *J Health Econ* 2005; **24**: 365–89.
- 44 Conibear L, Butt EW, Knot C, Arnold SR, Spracklen DV. Residential energy use emissions dominate health impacts from exposure to ambient particulate matter in India. *Nat Commun* 2018; **9**: 617.
- 45 McDuffie EE, Martin RV, Spadaro JV, et al. Source sector and fuel contributions to ambient PM<sub>2.5</sub> and attributable mortality across multiple spatial scales. *Nat Commun* 2021; **12**: 3594.
- 46 Meng W, Zhong Q, Chen Y, et al. Energy and air pollution benefits of household fuel policies in northern China. *Proc Natl Acad Sci USA* 2019; **116**: 16773–80.
- 47 Chowdhury S, Dey S, Guttikunda S, Pillarisetti A, Smith KR, Di Girolamo L. Indian annual ambient air quality standard is achievable by completely mitigating emissions from household sources. *Proc Natl Acad Sci USA* 2019; **116**: 10711–16.
- 48 Chatterjee D, McDuffie EE, Smith SJ, et al. Source contributions to fine particulate matter and attributable mortality in India and the surrounding region. *Environ Sci Technol* 2023; **57**: 10263–75.
- 49 Bond TC, Sun H. Can reducing black carbon emissions counteract global warming? *Environ Sci Technol* 2005; **39**: 5921–26.
- 50 Floess E, Grieshop A, Puzzolo E, et al. Scaling up gas and electric cooking in low- and middle-income countries: climate threat or mitigation strategy with co-benefits? *Environ Res Lett* 2023; **18**: 034010.
- 51 Emeterie M, Olaniyan I, Fadiji J, et al. Indoor air pollution: a review on the challenges in third world countries. *Air Soil Water Res* 2024; **17**: 11786221241239891.